arXiv:1302.5604v1 [nucl-th] 22 Feb 2013

Isospectral Bound State Potential from DDM3Y Effective Interaction

S. K. Dutta[1], D. Gupta[2], D. Das[2], and Swapan K. Saha[2]
[1] *Department. of Physics, B.G. College, Berhampore, Murshidabad 742101, India and*
[2] *Department. of Physics, Bose Institute, 93/1 A.P.C. Road, Kolkata 700009, India*

(Dated: February 25, 2013)

Theoretical investigation of $5/2^+$ resonance state of ^{11}Be was carried out using Supersymmetric Quantum Mechanics (SQM). The original two-body potential ($^{10}Be + n$) was constructed microscopically using a DDM3Y effective NN interaction. SQM converted this potential to an isospectral potential which is effective for detecting resonance states in the continuum. The resonance energy of the $5/2^+$ state is in good agreement with the experimental value.

PACS numbers: 21.45.-v, 27.10.+h

The theoretical procedure adopted in the present work, is appropriate to study resonances in weakly bound exotic systems [1–3]. Investigation of $5/2^+$ resonance state of ^{11}Be ($^{10}Be + n$) was undertaken. We started with a two-body potential which was constructed using density dependent M3Y (DDM3Y) effective NN interaction [4]. For the first time, the DDM3Y potential is used to study quasi-bound states of weakly bound nuclear system. The starting potential for our system presents a shallow well and a low, but very broad barrier. If the original potential $v(r)$ is strong enough to support a bound state of energy E where the normalized wave function $\Psi_0(r)$ satisfies the Schrodinger equation

$$[-\frac{\hbar^2}{2\mu}\frac{d^2}{dr^2} + v(r) - E]\Psi_0(r) = 0, \tag{1}$$

a straight forward calculation of resonance state from such potential faces numerical difficulties. So we use a well-known procedure in Supersymmetric Quantum Mechanics (SQM) to generate a one-parameter(λ) family of isospectral potentials from a given potential [5–7]. One parameter family of potentials $V(\lambda; r)$ are constructed by SQM from the original potential $v(r)$ by the following equation,

$$V(\lambda; r) = v(r) - 2\frac{\hbar^2}{2\mu}\frac{d^2}{dr^2}ln[I_0(r) + \lambda], \tag{2}$$

where

$$I_0(r) = \int_0^r [\Psi_0(r\prime)]^2 dr\prime. \tag{3}$$

The isospectral potentials thus generated look quite different from the original one but they present the same set of energy levels. Newly constructed isospectral potentials develop a deep and narrow well followed by a high barrier for small positive values of λ. The quasi-bound state in the original shallow potential will now be effectively trapped in the deep well of the isospectral potential. In this way SQM converts the original shallow potential to a deep-well isospectral potential, giving rise to enhanced trapping probability. This technique provides easy and accurate determination of energy and width of the resonant state. A free parameter λ is used to adjust the isospectral potential to desirable properties which enhances the trapping probability. Small values of λ enhances the probability of the system to be found within the potential well-barrier trap, resulting in sharp resonance peak in the trapping probability [2, 3]. For a resonance state having the same spin-parity of an existing bound state, isospectral potential is constructed from the ground state wave function. If there is no such bound state with the same spin parity J^π, we solve Schrodinger equation for a positive energy E subject to the boundary condition $\Psi_E(0) = 0$. The $\Psi_E(r)$ is normalized to a constant amplitude of oscillation in the asymptotic region. This $\Psi_E(r)$ is used for construction of isospectral potential.

In both cases isospectral potential approaches the original potential for $\lambda \to \infty$ and develops a deep and narrow well followed by a high barrier for $\lambda \to 0^+$ (Fig. 1). A judicious choice of λ is made, since for extremely small values of λ the numerical error in the evaluation of wave function increases. Probability of the system to be trapped within this enlarged well-barrier combination is measured by a function $C(E)$ given by

$$C(E) = \int_0^{r_B} [\hat{\Psi_E}(r\prime)]^2 dr\prime. \tag{4}$$

where $C(E)$ has a maximum at the resonance energy $E = E_R$. It is independent of the choice of λ. Width (Γ) of the resonance is obtained semi-classically from the mean life of the state using the time-energy uncertainty relation. The mean life is the reciprocal of the decay constant, which is expressed as a product of the number of hits on the barrier per unit time and the probability of tunneling through the barrier. The expression for (Γ) is given by

$$\Gamma = 2\sqrt{\frac{\hbar^2}{2\mu}}\frac{\exp(-2\int_b^c(\sqrt{\frac{2\mu}{\hbar^2}}(V(r;\lambda) - E_R)dr))}{\int_a^b \frac{dr}{\sqrt{(E_R - V(r;\lambda))}}}, \tag{5}$$

where a and b are the classical turning points of the well and b and c are the classical turning points of the barrier in $V(\lambda; r)$, corresponding to resonance energy E_R.



FIG. 1: One parameter family of isospectral potentials $V(\lambda; r)$ for $\lambda = 10^{-6}, 10^{-5}, 10^{-4}$ for the $5/2^{+}$ state of ^{11}Be



FIG. 2: Plot of probability $C(E)$ as a function of energy E for $\lambda = 10^{-6}, 10^{-5}, 10^{-4}$ for the $5/2^{+}$ state of ^{11}Be

By our theoretical procedure the resonance energy $E_R = 1.27$ MeV of ^{11}Be was found to be in good agreement with the experimental value ($E_{exp} = 1.274 \pm 0.018$ MeV) [8]. Resonance energy appears to be independent of the adjustable parameter λ (Fig. 2). Our original potential was slightly adjusted to reproduce the ground state of ^{11}Be. The width ($\Gamma = 124$ keV) slightly deviates from the experimental finding ($\Gamma_{exp} = 100 \pm 20$ keV) [8]. We also found that Γ is independent of λ. So we could conclude by saying that resonance energy calculated for $5/2^{+}$ resonant state of $^{11}Be(^{10}Be + n)$ agrees well with the experimental value. Although resonance width (Γ) is obtained semi classically, it agrees fairly well with the experimental result.

[1] S. K. Datta, T. K. Das, M. A. Khan, B. Chakrabarti, J. Phys. G: Nucl. Part. Phys. 29 (2003)2411
[2] S. K. Datta, T. K. Das, M. A. Khan, B. Chakrabarti, Few-Body Systems, 35 (2004)33
[3] S. K. Datta, T. K. Das, M. A. Khan, B. Chakrabarti, Intl. Jour. of Mod. Phys. E, 13 (2004) 811
[4] D. Gupta, E. Khan, Y. Blumenfeld, Nucl. Phys. A 773(2006)230; see references therein
[5] F. Cooper, A. Khare and U. Sukhatame, Phys. Rep. 251 (1995) 267
[6] A. Khare and U. Sukhatme, J. Phys. A22 (1989) 2847
[7] M. M. Nieto, Phys. Lett. 145 (1984) 208
[8] A. Shrivasta, Y. Blumenfeld, Phys. Lett. B 596 (2004) 54.